

**IMPORTANT NOTICE:** The following information is extracted from the label and is for information purposes only. When using the product, refer to the label which contains complete information regarding the use including applicable precautionary and first aid statements, etc.

## RESTRICTED USE PESTICIDE DUE TO HAZARD TO NONTARGET ORGANISMS

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

# P.C.Q.® PRO

ACTIVE INGREDIENT:  
Diphacinone (CAS #82-66-6): . . . . . 0.005%  
OTHER INGREDIENTS: . . . . . 99.995%  
TOTAL 100.000%

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions. Use only for sites, pests, and application methods described on this label.

Do not sell this product in individual containers holding less than 4 pounds of bait.

### USE RESTRICTIONS – For All Uses

Use only in rodent infested areas where exposure to agricultural crops will not occur. Do not contaminate food or food stuffs. Do not apply this bait at sites or to control pests not indicated on this label. Do not apply this product by application methods not specified in this label. Do not pile bait. Do not apply bait in or near actively growing food or feed crops or in areas where contaminating food or surfaces that may contact food is likely. Do not apply bait by aircraft at sites where bait may contaminate bodies of water. When applying bait along canals, and other waterways, use equipment that directs bait away from the water.



### NORWAY RATS, ROOF RATS AND HOUSE MICE



**IMPORTANT:** Do not expose children, pets, or nontarget animals to rodenticides. To prevent accidents:

1. Store unused product out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled, or unconsumed bait as specified on this label.

**SELECTION OF TREATMENT AREAS:** Determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

**USE RESTRICTIONS:** This product can only be used to control Norway rats, roof rats and house mice in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port of terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops.

Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations.

Tamper-resistant bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used if children, pets, nontarget mammals, or birds may access the bait.

In active rodent burrows, apply this product to control Norway rats, and roof rats within or beyond 100 feet of buildings and man-made structures, provided that infestations of these rodents have been confirmed. Bait must be placed no less than 6 inches into active Norway/roof rat burrows. Do not broadcast bait. Because Norway/roof rat infestations may occur in areas farther than 100 feet from buildings and man-made structures when the rodents have ample supplies of food and cover, efforts should be made to remove food trash, garbage, clutter and debris.

### APPLICATION DIRECTIONS:

**Norway and Roof Rats** - Apply 4 to 16 oz. of bait (usually at intervals of 15 to 30 feet) per placement in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease. Apply 1 to 2 oz. of bait deep into active rat burrows.

**House Mice:** Apply ¼ to ½ oz. (1 to 2 level tablespoons) of bait at 8 to 12 foot intervals in infested areas. Larger placements (up to 2 oz.) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

**FOLLOW-UP:** Replace contaminated or spoiled bait immediately. Using waterproof gloves, collect and dispose of all dead, exposed animals and leftover bait. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.



### CALIFORNIA GROUND SQUIRRELS

**USE RESTRICTIONS:** For control of California Ground Squirrels (*Spermophilus beechyi*) in industrial, commercial, and residential areas. Product must be used in a secured, tamper-resistant bait station or as a spot baiting application at the following sites: during the dormant season only for vineyards, fruit tree orchards (stone and pome), nut tree orchards, avocados, and citrus groves; rangeland, and pastures; noncrop borders of the listed use sites as follows: outside of fence rows, rights-of-way, canal banks, ditch banks, highways, levees, railroad lines and utilities. It may also be used in and around man-made structures such as agricultural buildings (but not inside of pens), homes, other permanent or temporary structures; campgrounds, golf courses, parks, athletic fields, recreational areas; ornamental plants; lawns; commercial horticultural nurseries, and plantations of forest trees.

Do not graze livestock or plant food or feed crops in spot-treated areas while bait is present. Applications in orchards, groves, and vineyards may only be made after harvest and during the dormant period and may not be made after tree and vine growth resumes in the spring.

Do not apply by aerial application. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers can be in the area during application.

**BAIT STATIONS:** Place secured tamper-resistant bait stations where California Ground Squirrel activity is present, usually near active burrows. Place bait stations up to 75 feet apart or near active burrows. Apply 2-4 lb. of bait per bait station. Bait stations should be checked every 3-5 days and filled as necessary until signs of bait consumption has ceased.

**SPOT BAITING:** Evenly scatter 1/4 cup in a measuring cup (0.1 lb.) of bait no larger than 40-50 square feet (i.e. a circular area no more than 8 feet in diameter), around the active California Ground Squirrel burrows and runways. Do not over-bait. Do not place bait in piles. Using the same procedure, make a second application 4 days after the first. Do not apply more than 10 pounds per acre per year.

**Surveillance and Follow-Up:** Dead ground squirrels should begin to appear in or near treated areas within 4 to 5 days after the first bait application. Visit the application site to monitor the effects of treatment 4 days after the first application and at subsequent intervals of 1 to 2 days. Wearing gloves, collect and properly dispose of visible carcasses of ground squirrels and other species found in or near treated areas. Dead animals buried on site must be buried a minimum of 18 inches below the ground surface, preferably deeper. Continue to collect and dispose of dead ground squirrels and search for non-target animals for at least 2 weeks after the last bait application, or longer if carcasses are still being found at that time. Carcass collection and burial should be performed in the late afternoon, near sundown, to reduce the potential for nocturnal, scavengers or predators to find carcasses on the ground surface. Wear gloves and use leakproof plastic bags or other suitable containers for transporting carcasses not buried on site. Where there is a continuous source of reinfestation, establish permanent bait stations and replenish bait as needed.



### MEADOW AND PINE VOLES

**INDUSTRIAL, COMMERCIAL, & RESIDENTIAL AREAS USE RESTRICTIONS:** For control of meadow voles (*Microtus pennsylvanicus*) and pine voles (*Microtus pinetorum*) in industrial, commercial, and residential areas. Areas include: in and around man-made structures such as agricultural buildings (do not apply in animal pens), homes & lawns (hand baiting into the burrow or tamper resistant bait station must be used for all residential uses and within 100 feet of a home), other permanent or temporary structures; campgrounds; golf courses; ornamental plants; commercial nurseries. If placed into a tamper-resistant bait station, may also be used in parks, athletic fields, and recreational areas.

Do not apply by aerial application. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers can be in the area during application.

**BURROW BAITING:** Identify surface trails and the mouths of holes leading to underground burrow systems. Place 1½ oz. (2 tablespoons) into each active burrow. Do not allow bait to be exposed on the surface. Do not disturb the runway system.

**BAIT STATIONS:** Place 2 lb. of bait in a tamper-resistant bait station in areas frequented by voles (near surface trails and burrows).

**GROUND BROADCAST BAITING:** Broadcast by hand or commercial spreader only within 100 feet of a man-made structure and only in commercial nurseries. Wear rubber gloves during hand-broadcast operations. Apply 10 lb. per acre. For high population areas, make a second application (10 lb. per acre) one to two months later. For pine vole application in states east of the Mississippi River, a rate of 20 lb. per acre may be required. Do not exceed 20 lb. per acre per calendar year.



### FRUIT TREE ORCHARDS (DORMANT)

**USE RESTRICTIONS:** For control of meadow voles (*Microtus pennsylvanicus*) and pine voles (*M. pinetorum*) in pome fruit and stone fruit tree orchards in the states listed below\*. Aerial application is only permitted in those following states where aerial is listed next to the state name. Apply only after fall harvest

(including drops), before new spring growth and when no rain or snow is expected for three consecutive days. Do not graze animals in treated areas. \*California, Connecticut, Georgia, Idaho (aerial), Massachusetts, Michigan (aerial), Missouri, Montana (aerial), North Carolina, New Hampshire, Ohio (aerial), Oregon (aerial), Pennsylvania (aerial), South Carolina, Utah (aerial), Virginia (aerial), Vermont (aerial), Washington (aerial), and West Virginia (aerial).

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers can be in the area during application.

**BURROW BAITING:** Identify surface trails and the mouths of holes leading to underground burrow systems. Place 1½ oz. (2 tablespoons) into each active burrow. Do not allow bait to be exposed on the surface. Do not disturb the runway system.

**BAIT STATIONS:** Place 2 lb. of bait in a tamper-resistant bait station in areas frequented by voles (near surface trails and burrows).

### BROADCAST BAITING:

Broadcast by hand, commercial spreader, or aerial (see use restrictions). Wear rubber gloves during hand-broadcast operations. Apply 10 lb. per acre. For high population areas, make a second application (10 lb. per acre) one to two months later.

# P.C.Q.® PRO



**BURROW BAIT FOR  
VOLES, CONTROL  
CALIFORNIA GROUND  
SQUIRRELS & MORE**



RESTRICTED-USE PESTICIDE



THE WORLD LEADER IN RODENT CONTROL TECHNOLOGY®

# P.C.Q. PRO



Durable pelleted bait formula



Burrow bait for voles in industrial, commercial & residential locations

## BROAD LABEL FOR USE AGAINST A WIDE VARIETY OF PESTS



Meadow & Pine Voles



California Ground Squirrels



Norway Rats, Roof Rats & House Mice

## PMP'S NEWEST WEAPON IN THE FIGHT AGAINST VOLES, GROUND SQUIRRELS & COMMENSAL RODENTS



- ▶ Restricted Use pesticide
- ▶ MEADOW & PINE VOLES: Approved for use in industrial, commercial & residential areas, including homes, lawns, golf courses and more. Burrow Baiting & Ground Broadcast Baiting applications allowed, as listed on the label. Expanded sites when used with a Bait Station.
- ▶ CALIFORNIA GROUND SQUIRRELS: Approved for use in Bait Stations or Spot Baiting Application in industrial, commercial and residential areas, including homes, lawns and more.
- ▶ Contains Diphacinone, a proven first-generation anticoagulant active ingredient
- ▶ Durable, weatherable pellets provide versatility for a variety of applications
- ▶ Highly palatable formula competes with natural food sources in the field
- ▶ Specially formulated to control California Ground Squirrels, Meadow & Pine Voles, Norway Rats, Roof Rats & House Mice
- ▶ Available in 12lb pails or 50lb bags

PRODUCT	CODE	CASE QTY
▶ PCQ PRO	PQ1012	12 lb. Pail
▶ PCQ PRO	PQ1050	50 lb. Bag




## THE WORLD LEADER

- ▶ The Largest Manufacturer of Rodenticides in the World
- ▶ Hold more patents than any other rodent control company in the world
- ▶ Hold more bait registrations of any rodent control company in the US
- ▶ Offer a complete line of rodent control products - pelleted and block bait, adhesive traps, mechanical traps and tamper-resistant bait stations



## CUTTING EDGE TECHNOLOGY

- ▶ Experts in rodenticide formulations
- ▶ Continuous product innovation
- ▶ Unmatched performance for more than 40 years



## OUTSTANDING PRODUCT

- ▶ Bell baits are proven in both lab and field settings, delivering results in a variety of conditions



THE WORLD LEADER IN RODENT CONTROL TECHNOLOGY®

www.belllabs.com | Madison, WI 53704 USA