PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals
Keep away from humans, domestic animals and pets, Harmful if swallowed, absorbed through the skin, inhaled, or in eyes, Do not breathe dust, Avoid contamination of feed or foodstuffs, Avoid contact with acids, Wash all utensils, spoons, or measuring devices thoroughly after using. Personal Protective Equipment (PPE)

All handlers, including anyone who applies bait and retrieves carcasses and unused bait, must wear

has been exempt by the FWS from the survey requirement

waterproof goves. Follow manufacturer's instructions for deaning/maintaining PPE, such as gloves. If no instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet.

ENVIRONMENTAL HAZARDS

This product is toxic to wildlife and fish. Birds and other wildlife feeding on treated bait may be killed. Do not apply where runoff is likely to occur. Do not apply this product directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

ENDANGERED SPECIES CONSIDERATIONS

coping Crane (Grus americana). Do not use this product during the period 30 days prior to arrival and 30 days after the whooping crane leaves known crane habitat. age aren une minopring craine reares known craine reactar. Aromu rabbit (Branhulague irlahoeneie). Do not use this product within one mile of alfalfa fields.

'ygmy ratort (*Beatrysigus* etamoenss). Uo not use this product within one mile of altital hebs. O not use this product within 7 km (4.34 miles) of any prairie dog town to finit inkis to the black-looted ferret Mustefalingingses from exposure to 2 miles plan of destruction of its prey base, unless the obtony is a, i an solated black-talled prairie dog town less than 80 acres in size; b) an isolated white-talled or Gunnison prairie log town less than 200 acres in size; c.) has been appropriately surveyed, using methods acceptable to th J.S. Fish and Wildlife Service, and found by the FWS not to be a suitable site for ferret reintroductions; c

UTAH: Utah Prairie Dog (Cynomys parvidens). Do not use this product in the range of th

user trains cog. CALFORNIA: Still mansh harvest mouse (Reithvodontomys raviventris), Do not use this product within 112 mile of salt marsh vogetation and/or brackish water wetlands which are located near or againent to San Pablo Bay and San Francisco Bay, Do not use this product within 112 mile of salt marsh vegetation and/or brackish water wetlands which are located in the Sacramento River below or adjacent to the contiliance of the cramento River and the San Joaquin River,

Morro Bay Kangaroo Rats (Dipodomys heermanni). Do not use this product within 2 1/2 miles of Bay Wood Park which is located on Morro Bay.

sey notion rain which is located on wonto bey. Tresno Kangaroo Rat (Dipodrtys nitratoides exitis), Follow the Interim Measures for Protecting Endangerei Species in the California Endangered Species Bulletin for the following California countries: Fresno, Kings

stephan's Kangaroo Rat (Dipodomys stephensi). Follow the Interim Measures for Protecting Endangere pecies in the California Endangered Species Bulletin for the following California coun on Bernardino, San Diego and Riverside.

utian Canada Goose (Branta canadensis feucopareia), Do not use this product from October to March in Federally closed goose hunting areas in the State of California. See current Duck and Goose Hunting

gainte answord a content inlay. Glant Kangaroo Rat (Olpodomys Ingens), Follow the Interim Measures for Protecting Endangered Species in the California Endangered Species Bulletin for the following California counties: Fresno, Kern, Kings, Merced

onterey, San Benito, San Luis Obispo, Santa Barbara and Tulare. nomerey, san Benno, san cuis cotepo, santa Barcaria and rulare. Injoin Kangareo Rat (Dipodomys nitratoides nitratoides). Follow the Interim Measures for Protectin Indangered Species in the California Endangered Species Bulletin for the following California counties: Ken

San Bernardino Kangaroo Rat (Dipodomys merriami parvus), Follow the Interim Measures for Protection dangered Species in the California Endangered Species Bulletin in the following California counti

San Bernarding and Riverside TEXAS: Attivater's greater prairie chicken (Tympanuchus cupido attivateri). Do not use this product in the As: number's greater plante criticien (ymparachia capado almatery, to induce insproduce and occupied range of the Affixades's praine chicken in the following Texas countiles: Afranasa, Aus oria, Colorado, DeWitt, Fort Bend, Galveston, Golad, Harris, Refugio, Victoria, Waller and Wharton.

PUERTO RICO: Yellow-shouldered Black Bird (Agetaius xanthomus) and Puerto Rican Plain Pigeo (Calumba incrnata wetmore). This product must not be used in the following areas except in tamper proof ba boxes: within 10 kilometers of Central Aquirer, Lago Cidra, Cebba, San German, 15 kilometers or e Esperanza, south of Highway 2 from city of Mayanuez to the city of Ponce and all Mona Island.

La esperienza, soum or ingrively 2 introncty or investigatez to the day or inchore and at montal station. OMICTEC: It is a food offense to use any specifición in a manner hat resulto in the debt of a member of an endangered species. The use of this product may pose a hazard to Federally designated endangered species deposes, Sugasteria messures to protect contagred species by nour county may vestigate in an EPA "finerim Measures" pampirtet, You may call EPA's Endangered Species Hotline, 1-800-447-3813, to: find out if an "Interim Measures" pamphlet exists for your county and have one sent to you. You can also consu your local county extension office or pesticide state lead agency to determine whether they have imposed an irements in your area to protect endangered and threatened species.

DIRECTIONS FOR USE

It is a violation of Federal betwo to see The INECTIONS FOR USE

READTHGL AREST, see The violation is makent reconsistent with its labeling.

READTHGL AREST, see The view the limit policy client is makent reconsistent with the labeling.

READTHGL AREST, see The view the limit policy client is reconsistent to the label.

Public Properties of the Violation of the labeling of the la

UNITATIONS

WINTATIONS

WINTAT

USE RESTRICTIONS: For control of Galfornia Ground Squirrels, (Spermophilus, beechey) on noncrop rights-of-way adjacent to canal and disch banks and other noncrop borders during tate spring and early summer. This product must not be applied or made, near residential areas, over water our greater paths are grown for the product must be applied or made, near residential areas, over water our greater paths are grown for the product must be applied or made, near residential areas, over water our greater paths are grown for the product of the pro

or feed.

PREBAITING: Prebaiting with 6 lbs, of untreated steamed crimped cats per acre one or two days prior to using

PREBATING: Prebating with 6 bits, of unfreated steamed cromped day garder one of the days prior to using 279 PRODENT OF \$27 PROSE OF \$

- RANGELAND AND AD JACENT NONCROP AREAS

PRAINE DOS
USE RESTRICTIONS: For control of black-lated prairie dogs (Cynomys Judovicianus), white-tailed prairie dogs
USE RESTRICTIONS: For control of black-lated prairie dogs (Cynomys Judovicianus), white-tailed prairie dogs
USE RESTRICTIONS: For control of black-lated prairie dogs
USE RESTRICTIONS
USE RESTRICTIONS
USE RESTRICTIONS
USE RESTRICTIONS
USE RESTRICTION
U Leucurus) and Gunnison (Zonn) pratine dogs (C. gunnison) on rangeland in the western United State
rith Dakota, South Dakota, Nebraska, Kansas, Oklahoma, Texas, New Mexico, Arizona, Colorado, Montane Idaho, Utah and Wyoming).
PREBAITING: Prebait with 4 grams (one teaspoon) of untreated steamed crimped cats per mound, one or two

days prior to baiting, to increase acceptance of treated baits by prairie dogs.

BAITING: Apply bait only after all or most of prebait is eaten and only to areas where untreated bait was

consumed. Establish observation period during prebaiting. Apply bait by hand as a six inch bait spot on edge of

each mound or in adjacent feeding areas. Application rate should not exceed 4 grams (one teaspoon) per bait spot. Treat during mid-summer through mid-winter period (July-January). Do not apply more than one bait application during this period. Dispose of spilled or urwanted bait by burial.

KANGAROO RATS TIONS: For control of the Ord's kangaroo rats (*Dipodomys ordil*), banner-tailed kangaroo rats pectabilis), and Merriam's kangaroo rats (*D. merriam*) in rangetand vegetation and noncrop areas. Do not this product within the occupied habitats of threatened or endangered species or subspecies of

cangaroo rats, HAND BATTING: Place one tablespoon (12 grams) of bait at 2 locations in feeding runs near the entrances to

___ MACADAMIA NUT ORCHARDS AND ADJACENT NONCROP AREAS _

moved from trees prior to shaker harvest or from ground prior to mechanized harvest, Orchards may be treated

up to 30 days prior to a harvesting round.

BURROW TREATMENT: Place 1-2 teaspoons (4-8 grams) of bait in each burrow. No preharvest interval is

required, provided nuts are not retrieved from burrows.

BROADCAST BARTING: Broadcast at the rate of five (5) pounds of bait per acre in orchard using a ground device or by hand. Aircraft may be used to treat adjacent noncrop areas if recommended and supervised by the Hawai Dept. of Health. The number of applications shall not exceed four (4) per year. The total amount of bait applied shall not exceed 20 pounds per acre per year.

ORCHARDS AND GROVES -

VOLES AND WHITE-FOOTED MICE
USE RESTRICTIONS: For control of meadow voles, printe voles, printe voles, Culfornia voles, long-tailed voles.
Design voles, montrol woles. Townsord or voles (Montral app) and white-footed mice in ordinaries and graves.

HAMD Bafffiles: Near the base of each intelested tree, place testaporofile quantities of that all 2-1 focations, either an undoor state or the mouth of robles leading to underground runner speers, Cover base afficially (ag., mats, bounds) or by pulling overhaining gasts back into place, Donat allow bast to be exposed on bure ground. Do not dischould be considered to the control of the state of the state

2-3 pounds per acre of infested trees, BROADCAST BAITING: Under infested trees broadcast bait eventy by cyclone seeder or by hand. Wear rubber

gloves during hand broadcast operations. Concentrate balting in the areas with the heaviest vegetative cover, Do not broadcast on bare ground or growing crops. Apply bait af rates from 6-10 lbs. per treated acre.

AERIAL APPLICATION: Broadcast bait at a rate of 6-10 lbs. per acre. Do not apply on bare ground or

GROUND SQUIRRELS
USE RESTRICTIONS: For control of California Ground Squirrels (Spermophilus beecheyl) and Richardson's Ground Squirrels (Spermophilus richardi) in orchards and groves, Apply bait after harvest while the orchard and

y prioses. Na with & lhe, of untreated steamed crimned nate ner acre one or two days prior to using

PREMATING: Printating with 6 lists of unthreaded element compared as per actio eits or into days print to using 250°PRODENT OFF 13 file kelps for himself and exceptation by California and Richardson's ground signified, HANDBARTING: Reply one feespoon if garries of hair or the ground near active burrows. HANDBARTING: Text of your conducting terminal replices, Discolated but using hand or ground-driven dispensing devices for to exceed 6 lbs, per acro. Wear nobler gloves during hand-incodactal operations, Dispose of essensible from a print of the print of

POCKET GOPHERS.
USE RESTRICTIONS: For control of plains pocket gophers (Geomys bursarius), southeastern pocket gophers (G. pinetis), yellow faced pocket gophers (Pappogeomys castanops) and pocket gophers (Thomomys spp.) in orchards and groves. Balt must be applied directly into pocket gopher's underground burrow systems or into

underground runnis and growes, but miss to expende entering improved general contents and contents of underground runnis and contents of the c ed resistance from soil.

due to decreased resistance from soil.

Applying Bait to Runways: Using a long-handled spoon or mechanical probe, drop one teaspoon (4 grams) of bait into the main runway at each based point, Cover the opening with sod, a rock, or soit to exclude light, Take pare not to cover bait with soit Depending upon found density make two to be dain placements part burrow system Recover and bury all spilled bait. Applied in his manner, one pound of bait will treat not eight acres, depending upon the number of pocket gopher burrow systems present.

BURROW BUILDER: Follow manufacturer's instructions for the type of equipment used. Calibrate equipment to

drop a teaspon quantity (4 grams) of bair at 4.5 foot intervals in the artificial burrow made by the at a rate of 2.3 lbs, of bair per acre, Pick up and bury all spilled bair.

SUGARCANE

USE RESTRICTIONS: For control of Polynesian rats (Ratus exulans), Norway rats (R. norvegicus), roof rats (R. rattus), rice rats (Cryzomiys palustris), Florida water rats (Meofiber alleni), and outton rats (Sigmodon hispidus)

in sugarcare felder.

PREBATING: Prebating with three pounds of untreated steamed critical operators one or two weeks prior to using 20° POCIDIT ON BMT All 70 may enhance ball acceptable by risk; to using 20° POCIDIT ON BMT All 70 may enhance ball acceptable by risk; and the rate of 5 bs, of ball per acceptable priority of the rate of 5 bs, of ball per acceptable priority of the rate of 5 bs, of ball per acceptable priority of the rate of 5 bs, of ball per acceptable priority of 50° Associated by the rate of 50 number of applications is 4 per crop or 20 lbs, per acre (0.4 b, ai/4) per crop, and the PHI is 90 days. For a 12-month cycle crop, the maximum number of applications is 2 per crop or 10 bs, per acre (0.2 lb, ai/4) per crop, and the PHI is 90 days. For all Crop cycles, the retreatment internal is 30 days.

NURSERY STOCK, CONIFER/CHRISTMASTREE, POPLAR/COTTONWOOD, ____ ORNAMENTAL AND NON-BEARING FRUIT TREE PLANTATIONS

USE RESTRICTIONS: For control of meadow voles, prairie voles, pine voles, Calfornia voles, Oregon voles,

USE RESTINATIONS: For control of meadow cycles, prains vietes, plan viete, Calibrian vietes, Drogon vietes, minimutal vietes and Tomescards vised (Microbia sep) in nursey stock, and conflictor/instruss tree, popularizonecod, ornamental, and non-bearing fruit tree plantations. Do not apply by air. MICROBIACHEST MICROBIACHEST VIETES (MICROBIACHEST VIETES VI

GRAPE VINEYARDS

USE RESTRICTIONS: For control of deer mice (Perconyesus maniculatus), white-footed mice (P. leucopus), ddfield mice (P. polianatus, except for threatened or endangered subspecies), meadow voles, prairie voles and pine voles Microtus spp.), and meadow jumping mice (Zapus hudsonius) in grape vinevards. Do not use this product in areas of Alabama and Florida that are occupied by threatened or endangered subspecies of the oldfield mouse.

BROADCAST BATTING: Broadcast bait evenly on the ground between the rows by cyclone seeder or by hand. Wear.

rubber gloves during hand-broadcast operations. Apply balt at a rate of 6-10 bs, per acre, Do not apply by air

RESTRICTED USE PESTICIDE DUE TO HAZARDS TO NONTARGET SPECIES

or retail sale and use only by Certified Applicators or persons under their direct upervision and only for those uses covered by the Certified Applicator's Certification.

PORT AG

OAT GRAIN BAIT FORMULATION

FOR CONTROL OF GROUND SQUIRRELS, PRAIRIE DOGS, RATS, VOLES, MICE, MEADOW MICE, DEER MICE, KANGAROO RATS, AND GOPHERS.

ACTIVE INGREDIENT:	
Zinc Phosphide: (CAS# 1314-84-7):	
INERT INGREDIENTS:	
	TOTAL 100%

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID

Have label with you when seeking treatment advice, including 1-877-854-2494.

Any persons applying zinc phosphide products and experiencing signs and symptoms such as nausea, abdominal pain, tightness in the chest, or weakness, should be see by a physician immediately.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice or transport

patient to the nearest hospital.

Do not administer anything by mouth or induce vomiting unless told to do so by the

poison control center or doctor. IF ON SKIN OR CLOTHING:

Take off contaminated clothing

Rinse skin immediately with plenty of water for 15-20 minutes.

 Call a poison control center or doctor immediately for treatment advice. IF INHALED:

Move person to fresh air.

If person is not breathing, call an ambulance, then give artificial respiration, preferably mouth to mouth, if possible,

Call a poison control center or doctor immediately for treatment advice. IF IN EVES

Hold eve open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing ever Call a poison control center or doctor immediately for treatment advice

See side panels for additional precautionary statements

NET WEIGHT: 45 lbs (20.41 kg)

EPA REG. NO. 12455-102 FPA FST NOs 12455-WI-1K 12455-WI-2P 12455-WI-3W Superscript is first letter of the lot number.





SQUIRREL OATS

- CROPI AND RANGEI AND AND NONCROP AREAS

USE RESTRICTIONS: FOR control of laten podest opches (Geomys bursanism), southeastern pocket gophers (G. pinelsk), yellow-faced pocket gophers (Papagagenmys castanops) and pocket gophers (Thimmomys gop.) in cropland (including alfalfa), rangeland, reforestation areas, Christmas trees, confleripopheriotrotinewood plantations and monorop areas (including lawns; goff courses and parisk), Balt must be applied directly into pocket gopher's underground burrow systems or into underground tunnels created by burrow-builder machines. Do no apply bait above ground or directly to crops grown for food or feed. Do not apply by use of burrow-builders in

HAND BAITING: Follow the directions indicated below to locate and treat pocket gophers' burrow systems. Locating Underground Rurways: Use a specially designed gopher probe, a metal rod, a strong smooth stick, or other suitable implement to probe the ground 10 to 15 inches away from fan shaped gopher mounds. Begin probing on the flat side of the fan. When the main runway has been entered, the probe will drop about 2 inches

que to decreaseu resisiance from soil.

<u>Applying Bait to Runways:</u> Using a long-handled spoon or mechanical probe, drop one teaspoon (4 crams) of bait <u>Anothern Set in Remense; Using a kny-handled spoon or menhandal probe, drop one lesspoon (a gramp) of the beginning through the beginning through the beginning through conf, or gold to exclude (e.g.f., Zales, and to cover betal with soil. Depending upon mounted density, make he to fine the able placements gate faitness age in the forecast enabling selection and coverage that the soil of the soil a faitness of the soil and placement and the soil and selection and selection and the soil and selection and selection and the selecti</u>

ALFALFA -USE RESTRICTIONS: For control of meadow voles, long-tailed voles, California voles, Oregon voles, r votes and Townsend's votes (Microtus spp.) and Richardson's ground squirrels (Spermophilus richardsonil) in altafa, All applications must occur shortly after a cutting of the hay, and/or prior to the next growth's attaining a length of 2 inches. Alfalfa forage from treated areas must not be harvested until it reaches maturity. This use is restricted to Montana, California, Idaho, Oregon, and the state of Washington.

PREBATTING: When bait is to be used to control Richardson's ground squirrels, prebaiting with untreated steamed crimped oats at a rate of 6 lbs, per acre 2-3 days before ZP® RODENT OAT BAIT AG is applied is likely

ablamed or myclo data is a raise of Ex., per a circ. 25 days a brotis x²⁶ HULBH LOA is 291 Aci a spite as leavy ABDOLOCATE BATH. The most circ. up to investable by air in grant-chronic dispersion for person, Aprily at a state of up to 10 bits, (C2. b. active importent) per a circ. More self-or on application privant I Rejainations or gound supervised are being particle. Hole are been impedied, a second bat large legication may be made left to days. A maximum of 20 bits, (Ic4. b. active importent) per a care may be applied per year. Do not apply by air when wind voicity actives 10 mile. Do not apply in pile or permit piles to be twent by the operations.

TIMOTHY AND TIMOTHY/ALFALFA MINTURES PRODUCED FOR HAY -USE RESTRICTION TO NOTION THAN APPLIED WITH URBS PRODUCED PLAY HAT USE RESTRICTION TO NOTION THAN APPLIED WITH URBS PRODUCED PLAY HAT USE RESTRICTION TO ANOTION THAN APPLIED WITH URBS PRODUCED PLAY HAT USES, and Townsend's vales (Microsot spin) in chiral with the play the cavely given the cavely general individual sea than the cave that the cave the cavel to be ready by the average given by the cavely given the plant than the cave that the cave that the cave the cavel the

Wear rubber gloves during hand-broadcast operations. A maximum of 2 applications by ground may be made at the rate of 5 to 10 lbs, per acre, one in the fall after the last cutting and one in the spring when timothy and timothy legume mixtures are still dormant. Do not apply in piles or permit piles to be formed by equipment.

TIMOTHY PRODUCED FOR SEED

USE RESTRICTIONS: For control of meadow vokes, long-tailed vokes, California vokes, Oregon vokes, mountain votes, and Townsend's votes (Microtius spp.) in timothy during the non-growing season. Op not apply by air, Do not apply to actively growing imothly or timothy/ailala or timothy/dover mictures. A minimum of 158 days must pass between an application of 12P® IRODENT OAT BAIT. As and any livestock foraging activity. This use is restricted

to Idaho, Oregon and the state of Washington.

BROADCAST BAITING: This product may be broadcast by hand, cyclone seeder, or ground driven dispensing consequence are two transportant may be broadcast by hand, cyclone seeder, or ground-driven dispensing devices at rates of up to 10 bs, (0.2 b, active ingredient) per acre during crop dominancy. Wear rubber gloves during hand-broadcast operations. A maximum of 20 bs, (0.4 lb, active ingredient) per acre may be applied annually.

USE RESTRICTIONS: For control of meadow votes, long-tailed votes, California votes, Oregon votes, mountain votes, Townsend's votes (Microtus spp.) and Richardson's ground squirrets (Spermophilis richardsonii) in groupseason and preglent applications in barler, Do not apply this product to barley within 50 days of harvest. This use is restricted to Montana, Idaho, Oregon and the state of Washington.

PREBATING: When bait is used to control Richardson's ground squirrets, prebaiting with untreated steamed crimped cats at a rate of 6 bs. per acre 2-3 days before ZP® RODENT OAT BAIT AG is applied is likely to enhance

BROADCAST BAITING: This product may be broadcast by air or ground-driven dispensing devices. Apply bait at a rate up to 6 bs, (0.12 b active ingredient) per acre. Make only one application per year if Richardson's ground squirrets are being targeted. If votes are being targeted, if votes are being targeted. If votes are being targeted per year. All

POTATOES -

USE RESTRICTIONS: For control of meadow vides, long-failed vides, California vides, Oregon vides, mountain vides, and Townsend's vides (Microtus spp.) in growing-season and preplant applications in potatoes. Do not apply this product to potatoes within 30 days of harvest. This use is restricted to Idaho, Oregon and the state

or reason space.

BROADCAST BAITING: This product may be broadcast by air or ground-driven dispensing devices. Apply bait once at a rate up to 10 lbs, (0,2 lb, active ingredient) per acre, or make two separate applications at rates of up to 5 bs, (0,1 lb, active ingredient) per acre. The maximum application rate per year is 10 lbs, (0,2 lb,

SUGAR BEETS -USE RESTRICTIONS: For control of meadow vols, long-tailed voltes, California voltes, Oregon voltes, mountain voltes and Townsend's voltes (Microtus spp.) in growing-season and preplant applications in sugar beets. Do not apply this product to sugar beets within 30 days of harvest. This use is restricted to Idaho, Oregon and the state

RROADCAST RATTING: This product may be broadcast by air or ground-driven dispension devices. Apply halt at rates up to 10 lbs. (0.2 lbs. active ingredient) per acre at planting. A second application may be made at the se rate. The maximum application rate per year is 20 lbs. (0.4 lb. active ingredient).

WHEAT USE ESTITETIVES For control of medicine of white EST white Tomardon is the Estimate of the EST white Tomardon is the Estimate of the EST white Tomardon is the Estimate of the EST point-present and product applications in wheat, Do not apply the product to wheat within 50 days of harvest. This was the restricted be Medicates (also control Rechardon's product or wheat within 50 days of harvest. This was the restricted the Medicates (also control Rechardon's proof squartes, probability with untreated steamed completed outs at an exist of the Estimate (also control Rechardon's proof squartes, probability with untreated steamed completed outs at an exist of the Estimate (also control Rechardon's proof squartes, probability with untreated steamed completed outs at an exist of Estimate (also control Rechardon's proof squartes, probability with untreated steamed complete outs at an exist of the Estimate (also control Rechardon's proof squartes, probability with untreated steamed proof section of the Estimate (also control Rechardon's proof squartes, probability with the control of the Estimate (also control Rechardon's proof squartes, probability with the control of the Estimate (also control Rechardon's proof squartes, probability with the Control Rechardon's proof squartes, probability with the Control Rechardon's proof squartes, probability with the control of the Control Rechardon's proof squartes, probability with the control of the Control Rechardon's proof squartes, probability with the Control Rechardon's proof squartes, probability with the control of the Control Rechardon's properties of the Control Rechardon's proof squartes, probability with the Control Rechardon's properties of the Cont

acceptance of the conclusion.

BROADCAST BAITING: This product may be broadcast by air or ground-driven dispensing devices. Apply bait at a rate of up to 6 fbs, (0.12 fb, active ingredient) per acre. Make only one application per year if Richardson's ground squirrets are being targeted. If votes are being targeted, a second application may be made within 25 days. The maximum application rate per year is 12 lbs. (0.24 lb. active ingredient) per acre.

see you are used to let the SEANS (DRY) because migrocartics part and the SEANS (DRY) because migrocartics part and the SEANS (DRY) because mice (false miscularly, doer mice (False miscularly doer mice) (False mice) and the state with an object to the protect by a state of the state

BROADCAST BAITING: This product may be broadcast by hand or ground-driven dispensing devices at a rate of 6 bs. (0.12 b. active) per acre. Wear rubber gloves during hand-broadcast operations. No more than one application per growing season is permitted,

RUSHRERRIES AND CANERERRIES -

USE RESTRICTIONS: For corroll of meditors, elect parts of the provider (Microsus spp.) in busiberies (highbard and lookush Substrems, current, electromics, possiberies and houseberies) and nonebush Electromics, current, electromics, possiberies and houseberies) and in carebories (blackbories), current, electromics, possiberies, possiberies,

interval is 70 days.

BROADCAST BAITING: This product may be broadcast by cyclone seeder, or by hand. Wear rubber gloves uring hand broadcast operations. When applying by hand, throw tablespoon amounts (12 grams) in cover along bushes, rock outcrops, and fence lines. Make up to 2 applications at a minimum interval of 21 days at the rate of 6 to 10 lbs, per acre per application.

at the text of \$1 to 10 b, ps and/or periphotation.

USE RESTRICTIONS' For control of praise viete (Monta) entropysism, mestow vides (M. ponsylvanicus), house mice (Mismarcus), destruction (Plancipus), thritterel-mind ground signarial, (Spirmachias) robitation manual signarial signarial (Spirmachias) robitation manual signarial signaria signaria

s during spills or equipment malfunctions,

Row Spaces Inches	4 lbs. / A Rate oz. Per 1000 ft.	5 lbs. / A Rate oz. Per 1000 ft.	6 lbs./ A Rate oz, Per 1000 ft.
20	2.5	3.1	3.7
30	3.7	4.6	5.5
36	4,4	5,5	6,6
38	4.7	5.8	6.9

- LAWNS, ORNAMENTALS, GOLF COURSES, PARKS AND NURSERIES -USE RESTRICTIONS: For control of meadow voles, prairie voles, pine voles, California voles, long-tailed voles
Oregon voles, mountain voles and Townsend's voles (Microtus spp.), Columbian ground squired. (Spermophilus columbianus), Richardson's ground squirrels (S. richardsonii), thirteen-lined ground squirrels (S. tridecern/ineatus), and California ground squirrels (S. beacheyl) in lawns, ornamentals, goti courses, parks,

and nurseries. Bait must not be applied on roads, over water or where plants are grown for food or feed, PREBATING: Prebail by placing one teaspoon (4 grams) of untreated steamed crimped oats around each active burrow opening or in each active rurway 2 or 3 days prior to using ZP® RODENT OAT BAIT AG to enhance

acceptance by ground squirrels or voles.

HAND BAITMG: Place 4 grams (one teaspoon) of bait around each active burrow opening or in each active nurwey, Allow bait to fall through to the ground surface but do not apply bait to bare ground. Do not place bait in heaps or place. nd squirrels or votes.

AROUND AGRICULTURAL AND INDUSTRIAL STRUCTURES -

USE RESTRICTIONS: This product may be used to control meadow voles (Microtus pennsylvanicus), prairie voles (M. cotrogaster). California voles (M. cotrogaster). California voles (M. cotrogaster). California voles (M. cotrogaster). The product may be used to control and industrial buildings, Do not apply bait inside

ced in tamper-resistant ball stations. These stations must be resistant to destruction by dogs and by children under 6 years of age and must be used in a manner that prevents such children from reaching into balt compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock,

raccoons, bears or other potentially destructive animals, or in areas prone to vandalism.

SELECTION OF TREATMENT AREAS: Determine where votes wit most likely find and consume the balt, Look

runways near buildings as well as signs of vote activity oriented toward the structures. PREBAITING: Prebailing with untreated steamed crimped cats may improve acceptance of this product and encourage vides to enter bail stations, Apply a teaspoor amount (4 grams) of prebail to each bail stations, Apply a te encourage valets to entre that stations, Apply a seaspoon amount (4 grams) of preshalt to each best station at testable 2 day before the scholar Station at testable 1 day before the scholar Station at testable 1 day before the scholar Station at testable 1 day to the scholar Station at the scholar Station Station at the scholar Station at the scholar Station at the scholar Station Station Station at internals of 2 the scholar Station at internals of 2 the scholar Station Station at internal scholar Station Station Station at internal scholar Station Station at internal scholar Station Stat

USE RESTRICTIONS: This product may be used at airports to control meadow voles (Microtus pennsylvanicus), prairie voles (M. ochrogaster), California voles (M. californicus), mountain voles (M. montanus), pine voles (M. pinetis), and white-flooted mice (Peromyscus leucopus). Do not apply bait to bare ground. Do not apply

PREBAITING: Prebaiting with untreated steamed crimped oats may improve acceptance of this product, Apply prebail at the same rate and using the same procedures that are to be used for applying the toxic ball. Apply prebail 2 or 3 days before the scheduled application of ZP® RODENT OAT BAIT AG.

HAND BAITING: Place a feaspoon (4 grams) of bait in surface runways or at the mouths of holes leading to

GROUND-BROADCAST BAITING: Broadcast ball evenly by hand, by cyclone seeder, or by mechanical spreader mounted on a tractor or 4-wheel drive all-terrain vehicle (ATV). Wear rubber gloves during hand-broadcast operations. Apply treated bait at rates from 6 to 10 lbs per acre. Do not broadcast bait on any area at an airport

STORAGE AND DISPOSAL

Do not continuint matter for for feet by strange of disposal.

STONAGES Stem of the feet of seed by strange of disposal.

STONAGES Stem of the feet of

gional Office for guidance. nearest EPA Hegional Uffice for guidance.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Dispose of empty container by placing in trash, at an approved waste disposal facility or by inci-

WARRANTY: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all of this material when such use and/or handling is contrary to label instructions.

AG5081