

POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONTRAC[®]

SOFT BAIT



ACTIVE CONSTITUENT: 0.05 g/kg BROMADIOLONE
Contains Denatonium Benzoate (Bittering Agent)

FOR THE CONTROL OF RATS AND MICE

PRIMARY PACK
CONTAINS 725 x 10 g SACHETS
WHICH ARE ILLEGAL TO SELL SEPARATELY.

DO NOT DESTROY OUTER CONTAINER WHILE PRODUCT STILL REMAINS.

NET 7.25 kg

Manufactured by:



Bell
LABORATORIES, INC.
www.belllabs.com
Windsor, Wisconsin 53598 U.S.A.
MADE IN USA

Australian Agent:
Bell Australia Pty Ltd
Level 50
120 Collins Street
Melbourne VIC 3000
Australia

APVMA Approval No.: 86331/120488

RESTRAINTS

DO NOT exceed 3 m between bait stations for mice, or 9 m for rats.
DO NOT place bait sachets in the open.
DO NOT use bait stations further than 2 m from buildings.

DIRECTIONS FOR USE

SITUATION	PEST	RATE	CRITICAL COMMENTS
In and around domestic homes, agricultural, industrial and commercial buildings, animal houses, farms, wharves, public service buildings, food factories, hospitals, inside transport vehicles (including ships) and around grain terminals.	Mice (<i>Mus musculus</i>)	1 to 4 soft baits per bait station or placement.	Eliminate as far as practicable, all alternative food sources. Bait areas where conditions might encourage rodent activity. Place baits so as to prevent access to non-target animals. Tamper resistant bait stations must be used at locations where access to non-target animals cannot otherwise be prevented. Inspect bait stations on a regular basis. Replace eaten or old/deteriorated baits. If eaten quickly, increase the number of bait stations. Continue observation and replenish until no more baits are taken. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.
	Rats (<i>Rattus norvegicus</i> , <i>Rattus rattus</i>)	2 to 20 soft baits per bait station or placement.	

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Determine areas where rats and/or mice will most likely find and consume bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places between floors and walls or in locations where rodents or their signs have been observed. Protect bait from rain or moisture. Replace contaminated or spoilt bait immediately. To prevent reinfestation, eliminate food, water and harbourage as much as possible. If reinfestation should occur, repeat treatment. Where a continuous source of reinfestation is present, establish permanent bait stations and replenish bait as needed. Collect and dispose of all dead animals and rodent faeces by burning or burial, and left over bait as per disposal instructions. Baiting for at least two weeks is necessary to reduce rat/mouse numbers.

PRECAUTIONS

DO NOT remove bait from sachet.

The use of this product for the control of protected native rodents requires permission from wildlife authorities.

DO NOT place bait in places accessible to children, domestic animals livestock and nontarget wildlife, unless in tamper resistant bait stations.

DO NOT place bait in areas where there is a possibility of contaminating food, livestock feed, or surfaces that come into direct contact with food. The product is restricted to use in an around buildings (within 2 metres) or enclosed spaces.

DO NOT apply bait to crops.

PROTECTIONS

PROTECTION OF WILDLIFE, FISH CRUSTACEANS AND ENVIRONMENT

DO NOT contaminate dams, rivers, streams or drains with the chemical or used container. Place baits in locations which are inaccessible to other animals and birds. This product and dead or affected rodents and rodent faeces are hazardous to pigs, cats, dogs, native wildlife and poultry.

PROTECTION OF LIVESTOCK

Bait, and dead or affected rodents are hazardous to birds, dogs, cats, pigs, poultry and non-target wildlife. Regularly search for, remove and destroy dead or affected rodents.

577AS-1

POISON

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

CONTRAC® SOFT BAIT

ACTIVE CONSTITUENT: 0.05 g/kg BROMADIOLONE
Contains Denatonium Benzoate (Bittering Agent)

FOR THE CONTROL OF RATS AND MICE

Contrac Soft bait is a highly palatable soft bait for the control of rats and mice including those resistant to other anticoagulants. For use in and around buildings.

Ready to use bait.
Kills in a single feed.
Soft Bait = palatable and attractive.

PRIMARY PACK
CONTAINS 725 x 10 g SACHETS
WHICH ARE ILLEGAL TO SELL SEPARATELY.

APVMA Approval No.: 86331/120488

NET 7.25 kg

CONDITIONS OF SALE

The use of CONTRAC® Soft Bait is beyond the control of the manufacturer. Therefore, no warranty, expressed or implied is given by Bell Laboratories, Inc. regarding its suitability or efficacy for any purpose. Bell Laboratories, Inc. cannot accept any responsibility for any consequence whatsoever resulting from the use of this product, save for those nonexcludable conditions implied by the Competition and Consumer Act 2010.



Australian Agent:
Bell Australia Pty Ltd
Level 50
120 Collins Street
Melbourne VIC 3000
Australia

STORAGE AND DISPOSAL

Store in the closed, original container in a well-ventilated area, as cool as possible. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food and feedstuffs, seed and fertilisers. Do not dispose of chemicals on site. Break, crush or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product. To the extent possible, untaken baits should be destroyed by burning or burial according to State or Territory requirements.

SAFETY DIRECTIONS

Poisonous if absorbed by skin contact or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with skin. When preparing product for use wear rubber gloves. If on skin and after each baiting, wash thoroughly with soap and water. After each day's use, wash gloves. Vitamin K1 (Phytomenadione) is antidotal.

FIRST AID INSTRUCTIONS

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26; New Zealand 0800 764 766. Vitamin K1 (Phytomenadione) is antidotal.

SDS

For further information please refer to the Safety Data Sheet which can be obtained from the supplier.

In a transport emergency dial 000, Police or Fire Brigade.

For specialist advice in an emergency only, call (02) 9037 2994 (24 h).

GHS Hazard and Precautionary Statements

May cause damage to organs through prolonged or repeated exposure.
Toxic to aquatic life with long lasting effects.

Do not get in eyes, on skin, or on clothing. Do not eat, drink or smoke when using this product. Avoid release to the environment. Use personal protective equipment as required. IF SWALLOWED: Rinse mouth. Do NOT induce vomiting. Collect spillage. Not combustible. Use extinguishing media suited to burning materials. Dispose of small quantities and empty containers by wrapping with paper and putting in garbage. For larger quantities, if recycling or reclaiming is not possible, use a commercial waste disposal service.

101620/09-20

Product Code: 919725

Batch No.:

D.O.M.



SPECIMEN LABEL



POISON
KEEP OUT OF REACH OF CHILDREN
CONTRAC® SOFT BAIT

ACTIVE CONSTITUENT: 0.05 g/kg BROMADIOLONE
Contains Denatonium Benzoate (Bittering Agent)

Before use read all directions on the outer pack.

APVMA Approval No.: 68331/11550 **NET 10 g**

**NOT TO BE
SOLD SEPARATELY**

2602AS