

WARNINGS

- Handle with care.
- Harmful when swallowed or absorbed through the skin.
- A mild eye irritant.
- Toxic to fish and other animals.
- Store in original container in a cool, dry place inaccessible to children and pets and away from food, feedstuffs and other chemicals.
- Keep containers tightly closed when not in use.
- Keep out of reach of children, uninformed persons and animals.
- In case of poisoning, call a doctor and present this label.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal climatic and storage conditions and the occurrence of resistance of the pests against the remedy concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS

- Avoid contact with skin.
- Wear protective gloves when handling bait.
- After use and in case of accidental skin contact, wash thoroughly with soap and water.
- Do not expose children, pets or non-target animals to rodenticide.
- Store unused product out of reach of children and pets.
- Apply bait in locations out of reach of children, pets, domestic animals and non-target wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtain bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas

• 587AF-1

open to hoofed livestock or other potentially destructive animals or in areas prone to vandalism.

- Do not contaminate ground water systems, surface water, ponds, waterways or ditches with rodenticide or its container.
- Destroy empty container by perforation and burn and do not re-use for any other purpose.
- Collect unused, spoiled or unconsumed bait and dead rodents and dispose of in a responsible manner.
- Prevent contamination of food, feedstuffs, eating utensils and drinking water.

SYMPTOMS OF POISONING

Nausea, vomiting, loss of appetite, extreme thirst, lethargy, diarrhoea, bleeding from gums and nose, bleeding under the skin, pallor, haemorrhage into and around elbow and knee joints, blood in urine and faeces.

FIRST AID TREATMENT

HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE

Eyes – Rinse eyes with plenty of clean water for at least 15 minutes.

Skin – Wash contaminated body area thoroughly with plenty of soap and water.

Ingestion – Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control centre or doctor.

NOTE TO PHYSICIAN

Administer Vitamin K₁ intravenously, 50-150 mg depending on prothrombin time on the first day. In severe cases, small transfusions of fresh blood can be considered.

NOTE TO VETERINARIAN

Administer Vitamin K₁ intramuscularly or orally as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

USE RESTRICTIONS

For control of Norway and roof rats and house mouse: Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food. This rodenticide must be applied in tamper-resistant bait stations. Do not broadcast bait.



Reg. No. L8259 Act No. 36 of 1947

A ready-for-use bait block for the control of roof and Norway rats and the house mouse in and around industrial, commercial and public buildings, in transport vehicles and in and around human dwellings and agricultural buildings.

Effective against warfarin resistant Norway rats and house mouse.

ACTIVE INGREDIENT:

Brodifacoum (coumarin anticoagulant). 0.05 g/kg

NET MASS 8 kg

Registered to:
MC Pharma (Pty) Ltd
Co. Reg. No.: 1999/009355/07
P.O. Box 336
Wierda Park 0149
Pretoria, SOUTH AFRICA
Tel. No.: +27 (0) 12 668 3019

Poison Information Centre Emergency Tel. No. 0861 555 777

Manufactured by:



Bell
LABORATORIES, INC. Madison, WI 53704 U.S.A.
www.belllabs.com
MADE IN USA

UN number 3808.99.00



**HARMFUL
SKADELIK**

Urban areas: This rodenticide may be used in and around the periphery of homes, industrial, commercial and public buildings. May also be used in transport vehicles (ships, trains, aircraft) and in and around related port or terminal buildings. May also be used in alleys.

Non-urban areas: This rodenticide may only be used in and around homes and agricultural buildings.

DIRECTIONS FOR USE: Use only as directed

GENERAL:

Norway rats and the house mouse may consume a lethal dose in one night's feeding with first dead rodents appearing four or five days after feeding begins.

SELECTION OF TREATMENT AREAS:

Determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain, direct sunlight and other extreme climatic conditions. Remove as much alternative food as possible.

APPLICATION DIRECTIONS:

MICE: Place 1 block per placement. Space placement at 2.5 to 3.5 metre intervals. Two blocks may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse activity cease.

RATS: Place 5 to 22 blocks (usually at intervals of 4.5 to 9.2 metres) per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

FOR USE IN SEWER: Thread wire through block and securely attach to a stationary structure such as the bottom step of a manhole ladder or a sewer grate. Allow just enough wire for the block to rest on manhole benching. If benching is not present, suspend block a few centimeters above the high

water mark or secure block on a board supported by opposing steps of the ladder. Securing the block in this manner will prevent removal by rats or water. Use at least 15 blocks per manhole. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead, exposed animals and leftover bait. To prevent re-infestation, limit sources of rodent food, water and harbourage as far as possible. If re-infestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

Each bait block in this container weighs nearly 10 or 20g, depending upon the presentation.

STORAGE AND DISPOSAL

Do not contaminate water, food or feedstuffs by storage or disposal.

Storage: Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

Pesticide disposal: In accordance with local and national regulations. Do not allow material to contaminate ground water system. Do not contaminate surface water.

Pesticide container: Do not re-use empty container. Do not contaminate ponds, waterways or ditches with chemical or used container. Destroy empty container by perforation and burn. If burning, stay out of smoke. Never use for any other purpose.

JAGUAR™ is a trademark of Bell Laboratories, Inc. U.S.A.

032816/03-16



Product Code: 933325

